

Panner's disease

Published on 05.12.2018

DOI: 10.1594/EURORAD/CASE.16265

ISSN: 1563-4086

Section: Musculoskeletal system

Area of Interest: Bones Extremities Musculoskeletal
bone Musculoskeletal joint

Procedure: Diagnostic procedure

Imaging Technique: Conventional radiography

Imaging Technique: MR

Imaging Technique: CT

Special Focus: Athletic injuries Case Type: Clinical
Cases

Authors: A. Anisau 1, 2, M.S. Posadzy 3, E. Mulier 1, F.
M. Vanhoenacker 1, 2, 4.

Patient: 7 years, male

Clinical History:

A 7-year old youth soccer player presented with periodic pain and slight swelling of his right elbow since two months, without previous trauma. There was a limited range of motion in the elbow.

Imaging Findings:

Conventional radiography showed joint effusion, irregular contours of the capitellum, increased bone density and a thin radiolucent line in the subchondral bone (Figures 1a and 1b). Follow-up with Cone Beam CT 2 months later depicted a crescent with vacuum phenomenon in the subchondral bone (Figures 2a and 2b), corresponding to the radiolucent line seen on radiographs.

Subsequent MRI revealed intact articular cartilage of the capitellum, predominantly low signal of the capitellum on both T1- and PD weighted images in keeping with sclerosis, and bone marrow edema in the adjacent humeral bone on FS PD weighted images (Figures 3a, 3b, 3c and 3d).

Discussion:

Panner's disease represents an osteochondrosis of the humeral capitellum. Osteochondroses affect epiphyses or epiphyseal equivalents and their progression ceases with skeletal maturity of the affected part [1]. Panner's disease occurs most frequently in boys under the age of 11 [2, 3].

Microtraumata from repetitive valgus stress and increased axial load are incriminated as the main pathogenetic factors, typically occurring in throwing sports and gymnastics respectively [2].

This may lead to disruption of vessels that supply the nucleus of the capitellar ossification centre, leading to ischemia, resulting in disturbed endochondral ossification [2].

Clinical presentation consists of pain, stiffness, swelling and a limited range of motion – especially an extension deficit, flexion deficit is less common [2]. The pain is activity-related [2].

Conventional radiographs may show an irregular contour, sclerosis, subchondral radiolucency, fragmentation and flattening of the capitellum [2, 3].

As the radiographical changes may be subtle, a low dose (Cone Beam) CT may be beneficial because of its greater sensitivity for detection of subtle changes and characterization of a subchondral vacuum phenomenon [4].

The radiolucent subchondral crescent represents gas accumulation in a subchondral fissure, which indicates osteonecrosis and impending articular collapse [4, 5]. This vacuum-phenomenon disappears when repair and re-

ossification occurs [4].

MRI shows heterogeneous signal intensity due to a variable degree of sclerosis, vacuum phenomenon and bone marrow edema in the ossified part of the capitellum, while the overlying cartilage is normal [2, 4]. The vacuum phenomenon has a similar low signal as the sclerotic changes and is therefore difficult to detect on MRI.

The majority of patients can be successfully treated conservatively, without serious morbidity [2].

The differential diagnosis of Panner's disease consists of osteochondrosis dissecans (OCD), which is seen in the second decade of life, when the ossification of the capitellum is complete [3, 5]. OCD affects both bone and overlying cartilage and may be complicated by intra-articular loose body formation and has an overall poorer outcome [2, 3]. MRI may detect cartilage defects, therefore it will be useful for distinguishing between Panner's disease and OCD.

Our patient was treated conservatively, with rest, avoidance of weight bearing and of leaning upon the right arm, promotion of passive movement and antiphlogistic medication. After six months there was a significant clinical improvement and a partial regression of radiological abnormalities.

Written informed patient consent for publication has been obtained.

Differential Diagnosis List: Panner's disease., Osteochondrosis dissecans., Ossification variant of the capitellum

Final Diagnosis: Panner's disease.

References:

- Danger F, Wasylw C, Varich L (2018) Osteochondroses. Semin Musculoskelet Radiol Vol. 22. (PMID:[29409078](#))
- Claessen FM, Louwerens JK, Doornberg JN, van Dijk CN, Eygendaal D, Bekerom MP (2015) Panner's disease: literature review and treatment recommendations. J Child Orthop 9(1): 9-17 (PMID: [25663360](#))
- Atanda Jr A, Shah SA, O'Brien K (2011) Osteochondrosis: common causes of pain in growing bones. Am fam physician 83(3) (PMID: [21302869](#))
- Anisau A, Posadzy M, Vanhoenacker F (2018) Panner's Disease: The Vacuum Phenomenon Revisited. Journal of the Belgian Society of Radiology 102(1)
- Resnick D (2002) Osteochondroses. Diagnosis of Bone and Joint Disorders, 4th ed., Chapter 74 3686-3741

Figure 1

a



Description: The capitellum has a slightly irregular articular contour (white arrow) and there is a radiolucent crescent in the subchondral bone (black arrows). Note slight joint effusion with displacement of the elbow fat pads (white asterisks). **Origin:** Image origin: Department of Radiology, AZ Sint-Maarten, Mechelen, Belgium

b



Description: The capitellum has a slightly irregular articular contour (white arrow) and there is a radiolucent crescent in the subchondral bone (black arrows). **Origin:** Image origin: Department of Radiology, AZ Sint-Maarten, Mechelen, Belgium

Figure 2

a



Description: Coronal image: The capitellum has an increased density and is slightly irregularly delineated (white arrow).

There is a subchondral crescent-shaped vacuum phenomenon (black arrow). **Origin:** Image origin: Department of Radiology, AZ Sint-Maarten, Mechelen, Belgium

b



Description: Sagittal image: The capitellum has an increased density and is slightly irregularly delineated (white arrow).

There is a subchondral crescent-shaped vacuum phenomenon (black arrow). **Origin:** Image origin: Department of Radiology, AZ Sint-Maarten, Mechelen, Belgium

Figure 3

a



Description: Coronal T1-weighted image: the capitellum is of heterogeneous signal intensity with areas of low signal anteriorly (white arrow).

The overlying articular cartilage is intact (small white arrowheads). **Origin:** Image origin: Department of Radiology, AZ Sint-Maarten, Mechelen, Belgium

b



Description: Sagittal T1-weighted images: the capitulum is of heterogeneous signal intensity with areas of low signal anteriorly and high signal posteriorly (white arrow). The overlying articular cartilage is intact (small white arrowheads). **Origin:** Image origin: Department of Radiology, AZ Sint-Maarten, Mechelen, Belgium

c



Description: Coronal FS PD images: the capitellum is of low signal (white arrow). The overlying cartilage is intact (white arrowheads). Note bone marrow edema in the distal humerus (black arrow).

Origin: Image origin: Department of Radiology, AZ Sint-Maarten, Mechelen, Belgium

d



Description: Sagittal FS PD images: the capitellum is of low signal (white arrow). The overlying cartilage is intact (white arrowheads). Joint effusion (black asterisks). Note bone marrow edema in the distal humerus (black arrow). **Origin:** Image origin: Department of Radiology, AZ Sint-Maarten, Mechelen, Belgium